

**OATS Allograft Protocol**

Name \_\_\_\_\_ Date \_\_\_\_\_

Procedure \_\_\_\_\_

Procedure Date \_\_\_\_\_

Frequency 1 2 3 4 5 times/week    Duration 1 2 3 4 5 6 weeks

\*\*\*Range of motion is an important progression of therapy, but limiting swelling is important.  
Respecting swelling will decrease pain and improve motion.\*\*\*

CPM: 2-3x a day for 2 hrs, start 0-40° increasing 5-10° per day as tolerated; goal 100° by 6 wks

	<b>Weight Bearing</b>	<b>Brace</b>	<b>ROM</b>	<b>Therapeutic Exercise</b>
<b>Phase I 0-6 weeks</b>	NWB	Week 0-1: brace locked in full extension Weeks 2-4: open brace by 20 degree increments DC brace once independent SLR without extensor lag	Initiate ROM	Gentle patellar mobs, AROM/PROM as tolerated, quad/hamstring sets, 4 way SLR, hamstring stretches  *Monitor incision *Ice/Elevate for swelling
<b>Phase II 6-8 weeks</b>	Partial weight bearing	None	Achieve ROM 0-130	ROM: prone hangs, flexinator/extensionator, stationary bike for ROM, PROM, scar mobilizations Strength: continue open chain quad/hamstring since PWB
<b>Phase III 8-12 weeks</b>	WBAT gradually	None	Full ROM	Initiate closed chain strengthening, wall squats, ¼ squats, hell/toe raises, proprioceptive activities, normalize gait

<b>Phase IV 3-6 months</b>	WBAT	None	Full ROM	Advance closed chain exercises and work on SLS activities, return to sport activities
--------------------------------	------	------	----------	---

**Comments:**

FCE \_\_\_\_\_ Work Conditioning/Work Hardening \_\_\_\_\_ Teach HEP \_\_\_\_\_

**Every patient's therapy progression will vary to a degree depending on many factors. Please use your best clinical judgment on advancing a patient. If other ideas are considered to improve patient's outcome do not hesitate to call.**

**Patient's recovery is a team approach: Patient, family/friend support, therapist, and surgeon. Every team member plays an important role in recovery.**

**Signature \_\_\_\_\_ Date \_\_\_\_\_**